

Direct U-Pb dating of Cretaceous and Paleocene dinosaur bones, San Juan Basin, New Mexico

James E. Fassett¹, Larry M. Heaman², and Antonio Simonetti³

¹Independent Research Geologist, 552 Los Nidos Drive, Santa Fe, New Mexico 87501, USA

²Department of Earth and Atmospheric Sciences, University of Alberta, Edmonton, Alberta T6G 2E3, Canada

³Department of Civil Engineering and Geological Sciences, Cushing Hall, Notre Dame University, Notre Dame, Indiana 46556, USA

Koenig et al. (2012) question two aspects of our paper (Fassett et al., 2011): (1) that we ignored a previously published critique by Lucas et al. (2009) that questioned the use of palynologic and paleomagnetic data to determine the age of the Ojo Alamo Sandstone, and (2) that we failed to provide sufficient detail regarding the dating methodology we employed. The first criticism is not relevant: our paper reporting the U-Pb dating of two dinosaur bones is independent of “previously published work.” (The Comment by Koenig et al. [2012] failed to mention that the previous critique by Lucas et al. [2009] of Fassett [2009a] was responded to in detail in Fassett, 2009b.)

The second criticism, that our dating methodology was not detailed, was due to the space limitations required by *Geology*; a more-detailed description of that methodology is in a manuscript in preparation by Heaman, Simonetti, and Fassett (Heaman, 2012, personal commun.). Koenig et al. note that there is scatter in the U-Pb data in excess of analytical uncertainty, especially for bone 22799-D. This occurred because there are regions in the bone affected by post-fossilization U-Pb disturbance, as fully discussed in the Heaman et al. manuscript in preparation (mentioned above).

Our analysis indicates, however, that there are regions in the bone samples where the geochemistry and U-Pb systematics are undisturbed. The enormous advantage of the U-Pb dating technique (Fassett et al., 2011) is that these pristine regions can be precisely targeted to obtain robust fossilization dates. Koenig et al. address the importance of establishing the time of bone fossilization because geochemical profiles in some fossil teeth and bone samples have shown that fossilization can take up to ~1 m.y., and is characterized by shallow geochemical gradients (Millard and Hedges, 1996; Trueman et al., 2004; Kohn, 2008; Koenig et al., 2009). Fossilization of most bone, however, is essentially complete in <100 k.y. (Trueman et al., 2004; Kohn, 2008). The geochemical gradients observed in bone BB-1 are steep and thus consistent with a short fossilization time. The Paleocene U-Pb age of 64.8 ± 0.9 Ma obtained for bone BB-1 is thus believed to represent the time of death within the quoted uncertainty.

The technique used to correct for common Pb in dating U-bearing minerals by laser ablation–inductively coupled plasma–mass spectrometry (LA-ICPMS) is important. We (2011) and other researchers have reported that apatite can contain a large proportion of common Pb; therefore, the accuracy of U-Pb dates in such cases is very sensitive to the isotopic composition assigned to the common Pb component. For LA-ICPMS analyses, it has been noted in numerous studies (Jackson et al., 2004; Storey et al., 2006; Chew et al., 2011) that it is not possible to make an accurate correction for common Pb based on measurement of the unradiogenic ²⁰⁴Pb content (i.e., the ²⁰⁴Pb method) because of isobaric interferences at mass 204 (in particular ²⁰⁴Hg). A U-Pb dating approach that does not require knowledge of the accurate abundance of ²⁰⁴Pb is the ²⁰⁷Pb method (Storey et al., 2006; Cox and Wilton, 2006; Simonetti et al., 2006; Chew et al., 2011). In this method, the common Pb isotopic composition is determined by pro-

jecting the uncorrected data on a Tera-Wasserburg diagram (²³⁸U/²⁰⁶Pb vs. ²⁰⁷Pb/²⁰⁶Pb), with the y-intercept delineating the ²⁰⁷Pb/²⁰⁶Pb ratio for common Pb. This common Pb composition can then be used to apply a common Pb correction to the measured ²⁰⁶Pb/²³⁸U ratios using well-established common Pb–radiogenic lead mixing equations (Chew et al., 2011). In studies where ICPMS and isotope dilution thermal ionization–mass spectrometry (ID-TIMS) analyses have been conducted on the same mineral containing high common Pb contents, this approach has yielded accurate dates (Aleinikoff et al., 2002; Storey et al., 2006; Simonetti et al., 2006; Simonetti et al., 2008; Chew et al., 2011). Thus, we adopted this well-established ²⁰⁷Pb method, and in our opinion, the U-Pb dates we reported are accurate. In contrast, Koenig et al. suggest that the ²³⁸U/²⁰⁶Pb isochron approach should have been used to determine ages because this method does not require any knowledge of the common Pb content or its isotopic composition. Although this method can be used for ID-TIMS dating of apatite, and in a crude way to assess the nature of isotopic disturbance in bone using ICPMS, it cannot be used to obtain accurate age results in conjunction with ICPMS techniques because of the large uncertainty associated with measurement of the combined ²⁰⁴(Pb+Hg) ion signal. This conclusion was reiterated in the recent study of Chew et al. (2011).

REFERENCES CITED

- Aleinikoff, J.N., Wintsch, R.P., Fanning, C.M., and Dorais, M.J., 2002, U-Pb geochronology of zircon and polygenetic titanite from the Glastonbury Complex, Connecticut, USA: An integrated SEM, EMPA, TIMS, and SHRIMP study: *Chemical Geology*, v. 188, p. 125–147, doi:10.1016/S0009-2541(02)00076-1.
- Chew, D.M., Sylvester, P.J., and Tubrett, M.N., 2011, U-Pb and Th-Pb dating of apatite by LA-ICPMS: *Chemical Geology*, v. 280, p. 200–216, doi:10.1016/j.chemgeo.2010.11.010.
- Cox, R.A., and Wilton, D.H.C., 2006, U-Pb dating of perovskite by LA-ICP-MS: An example from the Oka carbonatite, Quebec, Canada: *Chemical Geology*, v. 235, p. 21–32, doi:10.1016/j.chemgeo.2006.06.002.
- Fassett, J.E., 2009a, New geochronologic and stratigraphic evidence confirms the Paleocene age of the dinosaur-bearing Ojo Alamo Sandstone and Animas Formation in the San Juan Basin, New Mexico and Colorado: *Palaentologia Electronica*, v. 12, 146 p., http://palaentologia.org/splash/index12_1.html (last accessed January 2012).
- Fassett, J.E., 2009b, Response to critique by Lucas et al. (2009) of paper by Fassett (2009) documenting Paleocene dinosaurs in the San Juan Basin: *Palaentologia Electronica*, v. 12, no. 2, http://palaentologia.org/splash/index12_2.html (last accessed January 2012).
- Fassett, J.E., Heaman, L.M., and Simonetti, A., 2011, Direct U-Pb dating of Cretaceous and Paleocene dinosaur bones, San Juan Basin, New Mexico: *Geology*, v. 39, p. 159–162, doi:10.1130/G31466.1.
- Jackson, S.E., Pearson, N.J., Griffin, W.L., and Belousova, E.A., 2004, The application of laser ablation–inductively coupled plasma–mass spectrometry to in situ U-Pb zircon geochronology: *Chemical Geology*, v. 211, p. 47–69, doi:10.1016/j.chemgeo.2004.06.017.
- Koenig, A.E., Rogers, R.R., and Trueman, C.N., 2009, Visualizing fossilization using laser ablation–inductively coupled plasma–mass spectrometry maps of trace elements in Late Cretaceous bones: *Geology*, v. 37, p. 511–514, doi:10.1130/G25551A.1.
- Koenig, A.E., Lucas, S.G., Neymark, L.A., Heckert, A.B., Sullivan, R.M., Jasinski, S.E., and Fowler, D.W., 2012, Direct U-Pb dating of Cretaceous and Paleocene dinosaur bones, San Juan Basin, New Mexico: *Comment: Geology*, e253, doi:10.1130/G32154C.1.
- Kohn, M.J., 2008, Models of diffusion-limited uptake of trace elements in fossils and rates of fossilization: *Geochimica et Cosmochimica Acta*, v. 72, p. 3758–3770, doi:10.1016/j.gca.2008.05.045.
- Lucas, S.G., Sullivan, R.M., Cather, S.M., Jasinski, S.E., Fowler, D.W., Heckert, A.B., Spielmann, J.A., and Hunt, A.P., 2009, No definitive evidence of Paleocene dinosaurs in the San Juan Basin: *Palaentologia Electronica*, v. 12,

- 10 p., palaeo-electronica.org/2009_2/199/199.pdf (last accessed January 2012).
- Millard, A.R., and Hedges, R.E.M., 1996, A diffusion-adsorption model of uranium uptake by archaeological bone: *Geochimica et Cosmochimica Acta*, v. 60, p. 2139–2152, doi:10.1016/0016-7037(96)00050-6.
- Simonetti, A., Heaman, L.M., Chacko, T., and Banerjee, N., 2006, In-situ petrographic thin section U-Pb dating of zircon, monazite, and titanite using laser ablation-MC-ICP-MS: *International Journal of Mass Spectrometry*, v. 253, p. 87–97, doi:10.1016/j.ijms.2006.03.003.
- Simonetti, A., Heaman, L.M., and Chacko, T., 2008, Use of discrete-dynode secondary electron multipliers with Faradays—A “reduced volume” approach for in-situ U-Pb dating of accessory minerals within petrographic thin section by LA-MC-ICP-MS, *in* Sylvester, P., ed., *Laser Ablation ICP-MS in the Earth Sciences: Current Practices and Outstanding Issues: Mineralogical Association of Canada Short Course Series v. 40*, p. 241–264.
- Storey, C.D., Jeffries, T.E., and Smith, M., 2006, Common Pb-corrected laser ablation ICP-MS U-Pb systematics and geochronology of titanite: *Chemical Geology*, v. 227, p. 37–52, doi:10.1016/j.chemgeo.2005.09.003.
- Trueman, C.N., Behrensmeier, A.K., Tuross, N., and Weiner, S., 2004, Mineralogical and compositional changes in bones exposed on soil surfaces in Amboseli National Park, Kenya: Diagenetic mechanisms and the role of sediment pore fluids: *Journal of Archaeological Science*, v. 31, p. 721–739, doi:10.1016/j.jas.2003.11.003.