

Did Paleo-Tethyan anoxia kill arc magma fertility for porphyry copper formation?

Jeremy P. Richards^{1*} and A.M. Celâl Şengör^{2*}

¹Department of Earth and Atmospheric Sciences, University of Alberta, Edmonton, Alberta T6G 2E3, Canada

²Maden Fakültesi, Jeoloji Bölümü, and Avrasya Yerbilimleri Enstitüsü, İstanbul Teknik Üniversitesi, Ayazağa 34469 İstanbul, Turkey

ABSTRACT

The Tethyan orogen is host to numerous porphyry Cu ± Mo ± Au deposits, but the majority formed during subduction of the Neo-Tethyan ocean basin in the late Mesozoic–Cenozoic; very few deposits have been found associated with Paleo-Tethyan subduction. We propose that this sparsity is due to widespread anoxia in the Paleo-Tethyan ocean basin, leading to the generation of relatively reduced arc magmas that were infertile for porphyry Cu formation. A compilation of published geochemical data indicates that Neo-Tethyan arc rocks have higher average Cu contents and V/Sc and Sr/Y ratios compared to Paleo-Tethyan rocks, indicating higher magmatic oxidation states and greater fertility for ore formation during Neo-Tethyan subduction. Subduction of relatively reduced oceanic lithosphere, or reduction of normal moderately oxidized arc magmas by interaction with reduced lithosphere, can therefore destroy the ore-forming potential of arc magmatic suites.

INTRODUCTION

Phanerozoic arc magmas are generally hydrous and moderately oxidized, reflecting their derivation from partial melting of metasomatized suprasubduction zone asthenospheric mantle wedge material (Kelley and Cottrell, 2009; Zellmer et al., 2015). These magmas are fertile for the formation of porphyry Cu ± Mo ± Au deposits because chalcophile metals behave as incompatible elements in moderately oxidized magmas

[$\Delta\text{FMQ}+1$ to $+2$, where ΔFMQ (fayalite-magnetite-quartz) is the $\log f_{\text{O}_2}$ (oxygen fugacity) deviation from the FMQ buffer; Richards, 2015a]. Metals can thus be transported to upper crustal levels in magmas prior to partitioning into exsolving hydrothermal fluids, with reprecipitation in economic concentrations under favorable ore-forming conditions (Sillitoe, 2010).

The moderate oxidation state of Phanerozoic arc magmas is a critical metallogenic factor, because sulfur dissolved in the magma is mainly present as sulfate. In contrast, under more reducing conditions ($\Delta\text{FMQ} \leq 0$) the sulfur is predominantly present as sulfide, and early saturation in sulfide melts or solids will rapidly deplete the magma in chalcophile and siderophile elements (i.e., they will behave as compatible elements; Richards, 2015a), rendering the magma infertile for later magmatic-hydrothermal ore formation. It has been suggested that more reducing conditions may have characterized Precambrian subduction zones and arc magmas, thus explaining the rarity of Precambrian porphyry Cu deposits (Evans and Tomkins, 2011; Richards and Mumin, 2013). It is also possible that periods of deep ocean anoxia during the Phanerozoic Eon could have resulted in more reduced subduction zone conditions, similarly limiting magma fertility (Richards and Mumin, 2013).

An example of such a condition occurred in the Paleo-Tethyan ocean basin during the Permian Period, when almost complete isolation of the basin from global ocean circulation resulted in anoxic conditions and the deposition of thick sequences of organic-rich black shale (Figs. 1 and 2; Şengör and Atayman, 2009). Subduction of such sediments and




Figure 1. Distribution of igneous rocks associated with Paleo-Tethyan (Permian to Jurassic) and Neo-Tethyan (Cretaceous–Cenozoic) subduction and collisional closure (based on maps of Şengör et al., 1993; Richards, 2015b).

*E-mails: Jeremy.Richards@ualberta.ca; sengor@itu.edu.tr




Figure 2. Paleogeographic reconstruction of the Paleo-Tethyan ocean basin during the Late Permian (modified from Şengör and Atayman, 2009). Isolation from global oceanic circulation led to anoxia and the deposition of thick sequences of reduced seafloor sediments.

hydrothermally altered but unoxidized oceanic lithosphere would have generated unoxidized arc magmas ($\Delta FMQ \sim 0$) with low potential for porphyry ore formation. The lack of economic porphyry Cu deposits associated with Paleo-Tethyan arc magmatic systems could be explained by poor preservation of rocks of this age in the Tethyan orogen, and erosional loss of shallow porphyry deposits (generally formed <5 km below surface) (Wilkinson and Kesler, 2007). Alternatively (or additionally), they may simply not have formed due to the reduced nature of the subducting oceanic lithosphere.

Few reliable data exist for the oxidation state of Paleo-Tethyan arc rocks, many of which have undergone some degree of alteration or metamorphism. However, we present a compilation of published data for

least-altered Tethyan arc rocks that suggest that Paleo-Tethyan magmas were Cu poor compared with Neo-Tethyan magmas, possibly due to early sulfide saturation under more reducing conditions, rendering them less fertile for porphyry ore formation. These results support system-scale metallogenic models that require the optimal, efficient, and sequential operation of multiple processes for the eventual concentration of metals to form ore deposits (Richards, 2013). Inefficient or suboptimal functioning of any step can limit or destroy ore-forming potential, and reducing conditions in the subduction zone or deep lithosphere may be an early step that will render arc magmas infertile.

TETHYAN PORPHYRY Cu DEPOSITS

The Tethyan orogen is highly prospective for porphyry Cu \pm Mo \pm Au and associated epithermal Au \pm Cu deposits (Fig. 1), but the majority of known deposits are Cretaceous to Cenozoic in age, and are related to subduction and closure of the Neo-Tethyan ocean basin (Aghazadeh et al., 2015; Richards, 2015b). Examples include the Majdanpek porphyry Cu-Mo-Au deposit in Serbia (ca. 84 Ma), the Sar Cheshmeh porphyry Cu-Mo deposit in central Iran (ca. 14 Ma), and the Qulong porphyry Cu-Mo deposit in Tibet (ca. 16 Ma). The oldest known deposits occur in Tibet and the Indosinian porphyry belt of southwestern China, and include Yangla (ca. 233 Ma), Pulang (ca. 213 Ma), and Xietongmen (ca. 174 Ma). These Triassic–Jurassic systems are thought to be related to subduction and closure of various well-aerated Tethyan subbasins (Richards, 2015b). In contrast, there are no economic porphyry deposits known to be associated with subduction of normal Paleo-Tethyan oceanic lithosphere. This rarity of older deposits undoubtedly partly reflects their susceptibility to erosional loss, but some late Paleozoic–early Mesozoic upper crustal plutonic and volcanic rocks are preserved along the orogen (Fig. 1), suggesting that this cannot be the only explanation.

PERMIAN ANOXIA IN THE PALEO-TETHYS

Anoxia in the Paleo-Tethys ocean basin developed as a result of its equatorial location and restricted circulation from the late Carboniferous to Triassic (Fig. 2). The first evidence of reduced deep ocean conditions is found in late Carboniferous black shale sequences from Georgia (Dizi Series), northern and central Pamir (Kwahan, Bazardar suite), and north-eastern Tibet (Horpa Tso) (Fig. 3, columns 1–4; Şengör and Atayman, 2009).




Figure 3. Representative stratigraphic columns from the Paleo-Tethyan ocean basin during the Permian (modified from Şengör and Atayman, 2009). Isolation from global oceanic circulation led to anoxia and the deposition of thick sequences of reduced seafloor sediments.

1. Dizi series, Georgia
2. Kwahan section, northern Pamir
3. Bazardar suite, central Pamir
4. Horpa Tso, northwestern Tibet, China
5. Meishan, Zhejiang and Anhui provinces, China
6. Hanzhong-Guangyuan, Shaanxi-Sichuan, China
7. Xiejiaoguo and Huayun, Sichuan province, China
8. Eastern Alborz Mountains, northeast Iran
9. Paularo area, Carnic Alps
10. Brsnina section, Karawanken (southern Alps), Slovenia
11. Bükk Mountains, Hungary
12. Alanya Nappes, Taurus Mountains, southern Turkey
13. Qubu section, Tethyan Himalaya
14. Guryul Ravine, eastern Himalaya, Kashmir, India
15. Lampang, north Thailand

Anoxia eventually extended into shallow waters, causing bryozoan extinction on continental shelves starting in the Capitanian (Fig. 3, columns 5–15; Powers and Bottjer, 2007). In contrast, there is little evidence for anoxic conditions in the wider Panthalassa Ocean (Şengör and Atayman, 2009).

Northward subduction of the Paleo-Tethyan oceanic lithosphere below the Eurasian continental margin, and opening of the Neo-Tethyan ocean to the south by northward migration of the Cimmerian continental fragments (present-day central Turkey, Iran, and Tibet) effectively destroyed this anoxic ocean basin by the Early Triassic (Fig. 3; Şengör and Natal'in, 1996), limiting its lifespan to ~75 m.y.

TETHYAN ARC MAGMAS

The GEOROC database (Geochemistry of Rocks of the Oceans and Continents; <http://georoc.mpch-mainz.gwdg.de>) contains a large compilation of geochemical analyses of igneous rocks from the Tethyside orogenic belt. The vast majority of these analyses are of Cretaceous–Cenozoic Neo-Tethyan rocks from the Anatolian-Iranian section of the orogen. We have compiled separately data from early Mesozoic igneous rocks related to subduction and collisional closure of the Paleo-Tethyan ocean, reported from the Pontides, Greater Caucasus, central Iran, Turkmenistan, and Tibet. The data were filtered for lack of noted alteration and loss on ignition <2.0 wt%. (The data sets used are provided in Tables DR1 and DR2, with summary statistics in Table DR3, in the GSA Data Repository¹.)

The simplest assessment of magma fertility is the concentration of ore metals in fresh igneous rocks. Copper is routinely analyzed in geochemical packages, and a plot of Cu versus MgO as a measure of fractionation shows that the Paleo-Tethyan suite is consistently lower in Cu relative to Neo-Tethyan rocks (Fig. 4), suggesting a lower potential for ore formation.

Magmatic oxidation state is an important control on chalcophile element behavior, but this parameter is difficult to measure directly, and is commonly not reported. Instead, it has been suggested that trace element




Figure 4. Cu versus MgO in Paleo-Tethyan and Neo-Tethyan igneous rocks; median values are used to highlight the data trends [sources of data: GEOROC database (Geochemistry of Rocks of the Oceans and Continents; <http://georoc.mpch-mainz.gwdg.de>); Hess et al., 1995; Lemaire et al., 1997; Roger et al., 2003; Zhang et al., 2006a, 2006b, 2007; Cai et al., 2009, 2010; Dokuz et al., 2010; Karimpour et al., 2010; Yang et al., 2011, 2014; Mirnejad et al., 2013; de Sigoyer et al., 2014; Zhang et al., 2014; Xia et al., 2015; Zhang et al., 2016; Dai et al., 2017].

¹GSA Data Repository item 2017195, Tables DR1–DR3 (supplemental data and summary statistics), is available online at <http://www.geosociety.org/datarepository/2017/> or on request from editing@geosociety.org.

ratios such as V/Sc can be used as proxies for oxidation state (Li and Lee, 2004; Loucks, 2013). Li and Lee (2004) reported that depleted mantle-derived mid-oceanic ridge basalts (MORBs) with oxidation states close to $\Delta\text{FMQ} \sim 0$ have V/Sc ratios of 6.74 ± 1.11 , whereas Loucks (2013, 2014) suggested that oxidized igneous rocks associated with porphyry Cu deposit have V/Sc > 10 (at 58–70 wt% SiO_2). Selecting samples within this silica range, the Paleo-Tethyan suite has V/Sc ratios of 6.1 ± 2.2 (median = 5.6; n = 54), similar to MORB, whereas the Neo-Tethyan suite has elevated values of 9.1 ± 7.3 (median = 8.4; n = 447), closer to the range of fertile oxidized magmas proposed by Loucks (2013, 2014).

Loucks (2013, 2014) also proposed that fertile arc magmas have Sr/Y > 35, and $(\text{Eu}/\text{Eu}^*)/\text{Yb}_n > 2$ at 58–70 wt% SiO_2 [where $\text{Eu}^* = \sqrt{(\text{Sm}_n \times \text{Gd}_n)}$]. Higher values for these ratios reflect higher magmatic water content, which promotes early amphibole crystallization (preferentially removing Y and Yb from the melt relative to light rare earth elements) and suppresses plagioclase crystallization (delaying removal of Sr from the melt) (Richards and Kerrich, 2007), and may also correlate with arc oxidation state (Kelley and Cottrell, 2009). High magmatic water contents are obviously favorable for the formation of magmatic-hydrothermal ore deposits (Burnham, 1979). The Neo-Tethyan suite averages Sr/Y = 29.3 ± 23.0 (median = 26.0; n = 633) compared to lower values of 24.6 ± 18.9 (median = 18.1; n = 121) in the Paleo-Tethyan suite, whereas $(\text{Eu}_n/\text{Eu}^*)/\text{Yb}_n$ ratios are similar or marginally lower (Neo-Tethyan suite average = 1.57 ± 1.31 , median = 1.27; n = 323; Paleo-Tethyan suite average = 1.42 ± 0.77 , median = 1.35, n = 102) (see Supplementary Tables DR1–DR3).

Thus, the higher average Cu contents and V/Sc and Sr/Y ratios of the Neo-Tethyan suite indicate higher magmatic oxidation states and water contents, and suggest greater fertility for the formation of porphyry Cu deposits compared with the Paleo-Tethyan suite.

DISCUSSION AND CONCLUSIONS

Because porphyry Cu deposits require relatively oxidized, hydrous magmas to form, it has been proposed that their almost exclusive global restriction to Phanerozoic arc rocks reflects the delay of deep ocean oxidation, and therefore oxidative seafloor alteration and subduction zone metasomatism, until after the Neoproterozoic oxygenation event (Evans and Tomkins, 2011; Richards and Mumin, 2013). However, extensive periods of deep ocean anoxia have occurred during the Phanerozoic; one of the best documented cases is the Paleo-Tethyan ocean basin. Published data for igneous rocks formed in response to Paleo-Tethyan subduction suggest more reduced arc magmatic conditions and lower fertility for porphyry Cu formation, compared to the more typical, moderately oxidized conditions attending subduction of the Neo-Tethyan ocean basin, which resulted in widespread ore formation. Thus, the prospectivity of arc systems may reflect, as a primary control, the oxidation state of the subducted oceanic lithosphere: regions where reduced seafloor materials were subducted for extensive periods of time are unlikely to generate fertile arc magmatic systems. Elsewhere, secondary reduction of otherwise oxidized arc magmas may occur during ascent through crust containing reduced lithologies, resulting in early voluminous sulfide saturation and deep crustal loss of chalcophile metals, rendering the magmas infertile for later porphyry Cu formation (Tomkins et al., 2012; Hou et al., 2015).

ACKNOWLEDGMENTS

This contribution was supported by a Natural Sciences and Engineering Research Council of Canada Discovery Grant (RGPIN203099) to Richards. We thank Y. Zhang for providing unpublished geochemical data, R. Moritz for suggesting data sources, and R. Loucks for discussion of geochemical parameters. We also thank Katy Evans, Richard Sillitoe, and Steve Kesler for constructive reviews.

REFERENCES CITED

Aghazadeh, M., Hou, Z., Badrzadeh, Z., and Zhou, L., 2015, Temporal-spatial distribution and tectonic setting of porphyry copper deposits in Iran: Constraints from zircon U-Pb and molybdenite Re-Os geochronology: *Ore Geology Reviews*, v. 70, p. 385–406, doi:10.1016/j.oregeorev.2015.03.003.

- Burnham, C.W., 1979, Magmas and hydrothermal fluids, in Barnes, H.L., ed., *Geochemistry of hydrothermal ore deposits* (second edition): New York, John Wiley and Sons, p. 71–136.
- Cai, H., Zhang, H., and Xu, W., 2009, U-Pb zircon ages, geochemical and Sr-Nd-Hf isotopic compositions of granitoids in western Songpan-Garze fold belt: Petrogenesis and implication for tectonic evolution: *Journal of Earth Science*, v. 20, p. 681–698, doi:10.1007/s12583-009-0054-8.
- Cai, H., Zhang, H., Xu, W., Shi, Z., and Yuan, H., 2010, Petrogenesis of Indosinian volcanic rocks in Songpan-Garze fold belt of the northeastern Tibetan Plateau: New evidence for lithospheric delamination: *Science China Earth Sciences*, v. 53, p. 1316–1328, doi:10.1007/s11430-010-4033-9.
- Dai, L.-Q., Zheng, F., Zhao, Z.-F., and Zheng, Y.-F., 2017, Recycling of Paleotethyan oceanic crust: Geochemical record from postcollisional mafic igneous rocks in the Tongbai-Hong'an orogens: *Geological Society of America Bulletin*, v. 129, p. 179–192, doi:10.1130/B31461.1.
- de Sigoyer, J., Vanderhaeghe, O., Duchêne, S., and Billerot, A., 2014, Generation and emplacement of Triassic granitoids within the Songpan Ganze accretionary-orogenic wedge in a context of slab retreat accommodated by tear faulting, eastern Tibetan plateau, China: *Journal of Asian Earth Sciences*, v. 88, p. 192–216, doi:10.1016/j.jseas.2014.01.010.
- Dokuz, A., Karsli, O., Chen, B., and Uysal, I., 2010, Sources and petrogenesis of Jurassic granitoids in the Yusufeli area, northeastern Turkey: Implications for pre- and post-collisional lithospheric thinning of the eastern Pontides: *Tectonophysics*, v. 480, p. 259–279, doi:10.1016/j.tecto.2009.10.009.
- Evans, K.A., and Tomkins, A.G., 2011, The relationship between subduction zone redox budget and arc magma fertility: *Earth and Planetary Science Letters*, v. 308, p. 401–409, doi:10.1016/j.epsl.2011.06.009.
- Hess, J.C., Aretz, J., Gurbanov, A.G., Emmermann, R., and Lippolt, H.J., 1995, Subduction-related Jurassic andesites in the northern Great Caucasus: *Geologische Rundschau*, v. 84, p. 319–333, doi:10.1007/s005310050008.
- Hou, Z., Yang, Z., Lu, Y., Kemp, A., Zheng, Y., Li, Q., Tang, J., Yang, Z., and Duan, L., 2015, A genetic linkage between subduction- and collision-related porphyry Cu deposits in continental collision zones: *Geology*, v. 43, p. 247–250, doi:10.1130/G36362.1.
- Karimpour, M.H., Stern, C.R., and Farmer, G.L., 2010, Zircon U-Pb geochronology, Sr-Nd isotope analyses, and petrogenetic study of the Dehnow diorite and Kuhsangi granodiorite (Paleo-Tethys), NE Iran: *Journal of Asian Earth Sciences*, v. 37, p. 384–393, doi:10.1016/j.jseas.2009.11.001.
- Kelley, K.A., and Cottrell, E., 2009, Water and the oxidation state of subduction zone magmas: *Science*, v. 325, p. 605–607, doi:10.1126/science.1174156.
- Lemaire, M.M., Westphal, M., Gurevitch, E.L., Nazarov, K., Feinberg, H., and Pozzi, J.P., 1997, How far between Iran and Eurasia was the Turan plate during Triassic–Jurassic times?: *Geologie en Mijnbouw*, v. 76, p. 73–82, doi:10.1023/A:1003129028277.
- Li, Z.-X.A., and Lee, C.-T.A., 2004, The constancy of upper mantle f_{O_2} through time inferred from V/Sc ratios in basalts: *Earth and Planetary Science Letters*, v. 228, p. 483–493, doi:10.1016/j.epsl.2004.10.006.
- Loucks, R.R., 2013, Distinctive composition and genesis of copper ore-forming arc magmas: *Goldschmidt Conference 2013 Abstracts: Mineralogical Magazine*, v. 77, p. 1642.
- Loucks, R.R., 2014, Distinctive composition of copper-ore-forming arc magmas: *Australian Journal of Earth Sciences*, v. 61, p. 5–16, doi:10.1080/08120099.2013.865676.
- Mirnejad, H., Lalonde, A.E., Obeid, M., and Hassanzadeh, J., 2013, Geochemistry and petrogenesis of Mashhad granitoids: An insight into the geodynamic history of the Paleo-Tethys in northeast of Iran: *Lithos*, v. 170–171, p. 105–116, doi:10.1016/j.lithos.2013.03.003.
- Powers, C.M., and Bottjer, D.J., 2007, Bryozoan paleoecology indicates mid-Phanerozoic extinctions were the product of long-term environmental stress: *Geology*, v. 35, p. 995–998, doi:10.1130/G23858A.1.
- Richards, J.P., 2013, Giant ore deposits form by optimal alignments and combinations of geological processes: *Nature Geoscience*, v. 6, p. 911–916, doi:10.1038/ngeo1920.
- Richards, J.P., 2015a, The oxidation state, and sulfur and Cu contents of arc magmas: Implications for metallogeny: *Lithos*, v. 233, p. 27–45, doi:10.1016/j.lithos.2014.12.011.
- Richards, J.P., 2015b, Tectonic, magmatic, and metallogenic evolution of the Tethyan orogen: From subduction to collision: *Ore Geology Reviews*, v. 70, p. 323–345, doi:10.1016/j.oregeorev.2014.11.009.
- Richards, J.P., and Kerrich, R., 2007, Adakite-like rocks: Their diverse origins and questionable role in metallogenesis: *Economic Geology and the Bulletin of the Society of Economic Geologists*, v. 102, p. 537–576, doi:10.2113/gsecongeo.102.4.537.
- Richards, J.P., and Mumin, A.H., 2013, Magmatic-hydrothermal processes within an evolving Earth: Iron oxide-copper-gold and porphyry Cu \pm Mo \pm Au deposits: *Geology*, v. 41, p. 767–770, doi:10.1130/G34275.1.
- Roger, F., Arnaud, N., Gilder, S., Tapponnier, P., Jolivet, M., Brunel, M., Malavieille, J., Xu, Z., and Yang, J., 2003, Geochronological and geochemical constraints on Mesozoic suturing in east central Tibet: *Tectonics*, v. 22, 1037, doi:10.1029/2002TC001466.
- Şengör, A.M.C., and Atayman, S., 2009, The Permian extinction and the Tethys: An exercise in global geology: *Geological Society of America Special Paper* 448, 96 p., doi:10.1130/2009.2448.
- Şengör, A.M.C., and Natal'in, B.A., 1996, Palaeotectonics of Asia: Fragments of A Synthesis, in Yin, A., and Harrison, M., eds., *The Tectonic Evolution of Asia: Rubey Colloquium*: Cambridge, UK, Cambridge University Press, p. 486–640.
- Şengör, A.M.C., Cin, A., Rowley, D.B., and Nie, S.Y., 1993, Space-time patterns of magmatism along the Tethysides: A preliminary study: *Journal of Geology*, v. 101, p. 51–84, doi:10.1086/648196.
- Sillitoe, R.H., 2010, Porphyry copper systems: *Economic Geology and the Bulletin of the Society of Economic Geologists*, v. 105, p. 3–41, doi:10.2113/gsecongeo.105.1.3.
- Tomkins, A.G., Rebryna, K.C., Weinberg, R.F., and Schaefer, B.F., 2012, Magmatic sulfide formation by reduction of oxidized arc basalt: *Journal of Petrology*, v. 53, p. 1537–1567, doi:10.1093/petrology/egs025.
- Wilkinson, B.H., and Kesler, S.E., 2007, Tectonism and exhumation in convergent margin orogens: Insights from ore deposits: *Journal of Geology*, v. 115, p. 611–627, doi:10.1086/521606.
- Xia, R., Wang, C., Qing, M., Li, W., Carranza, E.J.M., Guo, X., Ge, L., and Zeng, G., 2015, Zircon U-Pb dating, geochemistry and Sr-Nd-Pb-Hf-O isotopes for the Nan'getan granodiorites and mafic microgranular enclaves in the East Kunlun Orogen: Record of closure of the Paleo-Tethys: *Lithos*, v. 234–235, p. 47–60, doi:10.1016/j.lithos.2015.07.018.
- Yang, T.N., Zhang, H.R., Liu, Y.X., Wang, Z.L., Song, Y.C., Yang, Z.S., Tian, S.H., Xie, H.Q., and Hou, K.J., 2011, Permo-Triassic arc magmatism in central Tibet: Evidence from zircon U-Pb geochronology, Hf isotopes, rare earth elements, and bulk geochemistry: *Chemical Geology*, v. 284, p. 270–282, doi:10.1016/j.chemgeo.2011.03.006.
- Yang, T.N., Ding, Y., Zhang, H.R., Fan, J.W., Liang, M.J., and Wang, X.H., 2014, Two-phase subduction and subsequent collision defines the Paleotethyan tectonics of the southeastern Tibetan Plateau: Evidence from zircon U-Pb dating, geochemistry, and structural geology of the Sanjiang orogenic belt, southwest China: *Geological Society of America Bulletin*, v. 126, p. 1654–1682, doi:10.1130/B30921.1.
- Zellmer, G.F., Edmonds, M., and Straub, S.M., 2015, Volatiles in subduction zone magmatism, in Zellmer, G.F., et al., eds., *The role of volatiles in the genesis, evolution and eruption of arc magmas*: *Geological Society of London Special Publication* 410, p. 1–17, doi:10.1144/SP410.13.
- Zhang, H.-F., Zhang, L., Harris, N., Jin, L.-L., and Yuan, H., 2006a, U-Pb zircon ages, geochemical and isotopic compositions of granitoids in Songpan-Garze fold belt, eastern Tibetan Plateau: Constraints on petrogenesis and tectonic evolution of the basement: *Contributions to Mineralogy and Petrology*, v. 152, p. 75–88, doi:10.1007/s00410-006-0095-2.
- Zhang, H.-F., Zhang, B.R., Harris, N., Zhang, L., Chen, Y.-L., Chen, N.-S., and Zhao, Z.-D., 2006b, U-Pb zircon SHRIMP ages, geochemical and Sr-Nd-Pb isotopic compositions of intrusive rocks from the Longshan-Tianshui area in the southeast corner of the Qilian orogenic belt, China: Constraints on petrogenesis and tectonic affinity: *Journal of Asian Earth Sciences*, v. 27, p. 751–764, doi:10.1016/j.jseas.2005.07.008.
- Zhang, H.-F., Parrish, R., Zhang, L., Xu, W.-C., Yuan, H.-L., Gao, S., and Crowley, Q.G., 2007, A-type granite and adakitic magmatism association in Songpan-Garze fold belt, eastern Tibetan Plateau: Implication for lithospheric delamination: *Lithos*, v. 97, p. 323–335, doi:10.1016/j.lithos.2007.01.002.
- Zhang, L.-Y., et al., 2014, Age and geochemistry of western Hoh-Xil-Songpan-Ganzi granitoids, northern Tibet: Implications for the Mesozoic closure of the Paleo-Tethys ocean: *Lithos*, v. 19, p. 328–348, doi:10.1016/j.lithos.2013.12.019.
- Zhang, Y., Niu, Y., Hu, Y., Liu, J., Ye, L., Kong, J., and Duan, M., 2016, The syn-collisional granitoid magmatism and continental crust growth in the West Kunlun Orogen, China—Evidence from geochronology and geochemistry of the Arkarz pluton: *Lithos*, v. 245, p. 191–204, doi:10.1016/j.lithos.2015.05.007.

Manuscript received 12 January 2017

Revised manuscript received 2 March 2017

Manuscript accepted 3 March 2017

Printed in USA