

## How local crustal properties influence the amount of denudation derived from low-temperature thermochronometry

**Rob Westaway**

School of Engineering, University of Glasgow, Glasgow G12 8QQ, UK

Luszczak et al. (2017) apply apatite fission-track thermochronometry (AFTT) to the Cenozoic landscape development of northern Britain, including the Lake District (LD) of northwest England. Their main conclusion is that previous workers (e.g., Green, 2002; Green et al., 2012) have overestimated denudation by overestimating the thermal conductivity ( $k$ ) of the rock column lost. However, Luszczak et al. infer much more denudation than these other studies, although with much lower Late Cenozoic uplift rates, contradictions that require comment. Luszczak et al. also dispute the inference by other workers of a higher-than-present paleo-geothermal gradient,  $\nabla T_p$ , caused by transient heating of the shallow crust during the Paleocene British Tertiary Igneous Province (BTIP) magmatism.

Luszczak et al. infer that the rock lost from the LD was Mesozoic mudstone and Chalk with  $k = 1.2\text{--}2.0$  W/m $^{\circ}\text{C}$  (cf. Green, 2002). Present outcrop in the LD is mostly Ordovician Borrowdale Volcanic Group (BVG), with offlapping Carboniferous and Permian-Triassic sediments (e.g., Holliday, 1993). For the BVG rocks, sampled by Green (2002),  $k$  is  $2.51 \pm 0.19$  W/m $^{\circ}\text{C}$  ( $\pm 1s$ ; Gunn et al., 2005). With a harmonic mean  $k$  of 1.6 instead of  $\sim 2.5$  W/m $^{\circ}\text{C}$ , the Cenozoic denudation estimated from Green's AFTT adjusts from  $\sim 700$  to  $\sim 450$  m. His arguments, and those by Green et al. (2012) and Westaway (2017), would not be significantly affected by this, but his original  $\sim 700$  m value agrees better with Holliday's (1993) geological estimate of  $\sim 700\text{--}1750$  m of denudation.

Luszczak et al. partitioned their preferred 2250 m of Cenozoic uplift in the LD with 1650 m during 62–57 Ma, at 0.33 mm/yr, followed by  $\sim 500$  m at 0.0088 mm/yr. However, uplands in northern England have maintained uplift rates of  $\sim 0.15\text{--}0.2$  mm/yr since the Early Pleistocene, with uplift by  $\sim 150\text{--}200$  m on this time scale and by  $\sim 300\text{--}400$  m since the Mid Pliocene (e.g., Waltham, 2013; Westaway, 2017). These relatively high rates, established independently of AFTT, are evidently unrelated to the BTIP magmatism. The uplift history preferred by Luszczak et al. is consistent with  $\sim 1250$  m of denudation over the highest part of the LD (978 m above sea level [a.s.l.]), far above Green's (2002) estimate. However, they did not report any analyses with  $<2000$  m of uplift assumed, and give no compelling reason for choosing their preferred solution, raising the possibility of significant overestimation. This is important, because 2250 m is roughly the uplift expected from isostatic modeling, after Brodie and White (1994), following the intrusion of  $\sim 5$ -km-thick underplating, as is observed (e.g., Barton, 1992). However, the Brodie and White modeling method is considered over-simplistic (e.g., Green et al., 2012; Westaway, 2017). As Luszczak et al. present no analyses with  $\sim 450\text{--}700$  m of denudation across the LD, their paper sheds no light on this key issue.

AFTT indicates that boreholes offshore of Britain record a 'pulse' of heating by the BTIP magmatism (e.g., Clift and Turner, 1998). Green's (2002) sampling of the BVG rocks up to 966 m a.s.l. in the LD provides an analogous vertical transect, with a  $\nabla T_p$  of 61  $^{\circ}\text{C}/\text{km}$ , indicating paleo-heat-flow of 61  $^{\circ}\text{C}/\text{km} \times 2.51$  W/m $^{\circ}\text{C}$ , or  $\sim 153$  mW/m $^2$ ,  $\sim 60\%$  above present, consistent with the offshore context. Green's analysis is not refuted by Luszczak et al., whose sites are all too low (a.s.l.) to resolve  $\nabla T_p$ . Luszczak et al. also argue that the Green et al. explanation for the increased  $\nabla T_p$  is wrong, stating "deep-seated magmatism has a negligible impact on the thermal structure of shallow crust" and citing Brown et al. (1994). However, Brown et al. assumed instantaneous underplating at the base of the crust, whereas the BTIP magmatism spanned millions of years (e.g., 61–52 Ma; Kent and Fitton, 2000). Following Clift (1999), many workers have realized that offshore thermal histories reflect

prolonged underplating by many small intrusions. Green et al. (2012) thus assumed that the top of the underplate remained at 1100  $^{\circ}\text{C}$ , the freezing point of mafic magma, for up to 10 m.y., to mimic this scenario. The resulting calculated shallow geothermal gradient thus increased by  $\sim 60\%$  within  $\sim 5$  m.y. of the start of the magmatism, explaining the AFTT evidence. The assumption of instantaneous magmatism evidently led Luszczak et al. to a mistaken conclusion.

Luszczak et al. rightly emphasize the importance of basing AFTT modeling on site-specific, rather than nominal, data. Nonetheless (as for any modeling), judgment is required as to which details are key. Apart from the higher-than-present  $\nabla T_p$ , their analysis also omits other aspects, including surface temperature fall (cf. Green et al., 2012), and causes of Cenozoic denudation and uplift other than the BTIP magmatism (cf. Green et al., 2012; Westaway, 2017). Aspects such as these, omitted from their modeling, have evidently led them to overestimate Cenozoic denudation; their analysis is thus not a sound basis for superseding the interpretation (e.g., Green, 2002; Green et al., 2012) of only modest ( $\sim 700$  m) Cenozoic denudation in uplands of northern England.

### REFERENCES CITED

- Barton, P.J., 1992, LISPB revisited: A new look under the Caledonides of northern Britain: *Geophysical Journal International*, v. 110, p. 371–391, <https://doi.org/10.1111/j.1365-246X.1992.tb00881.x>.
- Brodie, J., and White, N., 1994, Sedimentary basin inversion caused by igneous underplating: Northwest European continental shelf: *Geology*, v. 22, p. 147–150, [https://doi.org/10.1130/0091-7613\(1994\)022<0147:SBICBI>2.3.CO;2](https://doi.org/10.1130/0091-7613(1994)022<0147:SBICBI>2.3.CO;2).
- Brown, R., Gallagher, K., and Duane, M., 1994, A quantitative assessment of the effects of magmatism on the thermal history of the Karoo sedimentary sequence: *Journal of African Earth Sciences*, v. 18, p. 227–243, [https://doi.org/10.1016/0899-5362\(94\)90007-8](https://doi.org/10.1016/0899-5362(94)90007-8).
- Clift, P.D., 1999, The thermal impact of Paleocene magmatic underplating in the Faeroe–Shetland–Rockall region, in Fleet, A.J., and Boldy, S.A.R., eds., *Petroleum Geology of Northwest Europe*: Geological Society of London Petroleum Geology Conference Series, v. 5, p. 585–593.
- Clift, P.D., and Turner, J., 1998, Paleogene igneous underplating and subsidence anomalies in the Rockall–Faeroe–Shetland area: *Marine and Petroleum Geology*, v. 15, p. 223–243, [https://doi.org/10.1016/S0264-8172\(97\)00056-1](https://doi.org/10.1016/S0264-8172(97)00056-1).
- Green, P.F., 2002, Early Tertiary paleo-thermal effects in northern England: Reconciling results from apatite fission track analysis with geological evidence: *Tectonophysics*, v. 349, p. 131–144, [https://doi.org/10.1016/S0040-1951\(02\)00050-1](https://doi.org/10.1016/S0040-1951(02)00050-1).
- Green, P.F., Westaway, R., Manning, D.A.C., and Younger, P.L., 2012, Cenozoic cooling and denudation in the North Pennines (northern England, UK) constrained by apatite fission-track analysis of cuttings from the Eastgate Borehole: *Proceedings of the Geologists' Association*, v. 123, p. 450–463.
- Gunn, D.A., Jones, L.D., Raines, M.G., Entwisle, D.C., and Hobbs, P.R.N., 2005, Laboratory measurement and correction of thermal properties for application to the rock mass: *Geotechnical and Geological Engineering*, v. 23, p. 773–791, <https://doi.org/10.1007/s10706-003-3156-6>.
- Holliday, D.W., 1993, Mesozoic cover over northern England: interpretation of apatite fission track data: *Journal of the Geological Society*, v. 150, p. 657–660, <https://doi.org/10.1144/gsjgs.150.4.0657>.
- Kent, R.W., and Fitton, J.G., 2000, Mantle sources and melting dynamics in the British Palaeogene Igneous Province: *Journal of Petrology*, v. 41, p. 1023–1040, <https://doi.org/10.1093/petrology/41.7.1023>.
- Luszczak, K., Persano, C., Braun, J., and Stuart, F.M., 2017, How local crustal properties influence the amount of denudation derived from low-temperature thermochronometry: *Geology*, v. 45, p. 779–782, <https://doi.org/10.1130/G39036.1>.
- Waltham, T.C., 2013, Karst geomorphology, in Waltham T.C., and Lowe, D., eds., *Caves and Karst of the Yorkshire Dales*, vol. 1: Buxton, UK, British Cave Research Association, p. 65–92.
- Westaway, R., 2017, Isostatic compensation of Quaternary vertical crustal motions: coupling between uplift of Britain and subsidence beneath the North Sea: *Journal of Quaternary Science*, v. 32, p. 169–182, <https://doi.org/10.1002/jqs.2832>.