

# Multisourced metals enriched by magmatic-hydrothermal fluids in stratabound deposits of the Middle–Lower Yangtze River metallogenic belt, China

Yang Li<sup>1,2,3\*</sup>, David Selby<sup>1,4</sup>, Xian-Hua Li<sup>2</sup>, and Chris J. Ottley<sup>1</sup>

<sup>1</sup>Department of Earth Sciences, Durham University, DH1 3LE Durham, UK

<sup>2</sup>State Key Laboratory of Lithospheric Evolution, Institute of Geology and Geophysics, Chinese Academy of Sciences, 10029 Beijing, China

<sup>3</sup>Department of Geology and Geophysics, Yale University, New Haven, Connecticut 06511, USA

<sup>4</sup>State Key Laboratory of Geological Processes and Mineral Resources, School of Earth Resources, China University of Geosciences, Wuhan, 430074 Hubei, China

## ABSTRACT

**Stratabound deposits within late Carboniferous carbonate units in the Middle–Lower Yangtze River metallogenic belt are important copper producers in China. Hitherto, the genesis of these deposits has been debated, due to poor constraints regarding the timing and source of the mineralization. Proposed models include a late Carboniferous seafloor exhalative formation (SEDEX), or an Early Cretaceous magmatic-hydrothermal origin. These models imply different metal sources (basinal vs. magmatic fluid, respectively) and would require different exploration strategies. New pyrite Re-Os and trace-element results from the representative Xinqiao Cu-S-Fe-Au deposit favor a Cretaceous (ca. 138 Ma) magmatic-hydrothermal genesis over a SEDEX origin. The distinct initial <sup>187</sup>Os/<sup>188</sup>Os compositions ( $Os_i$ ) of different pyrite types (colloform  $Os_i = 1.35$  and euhedral grains  $Os_i = 0.79$ ), coupled with the pyrite trace-element abundance, indicate that the Os, and by inference other metals (e.g., Cu, Ag, Au), was sourced from a Cretaceous magmatic-hydrothermal system ( $Os_i = 0.74$ ) and Late Permian metalliferous black shales ( $Os_i = 7.56 \pm 3.76$ ). In addition, the genesis of Au-bearing stockwork pyrite veins hosted by the Carboniferous sandstone is best explained by the leaching of existing mineralization (e.g., porphyry Au-Mo) by Early Cretaceous magmatic-hydrothermal fluids. This is implied by the lack of common Os, high Re abundances (0.1–3.7 ppm), and highly variable Re-Os model ages (379 and 173 Ma), which are positively correlated with Re and total abundances of Co, Ni, Ag, Au, Tl, and Ba. This study highlights the importance of recycling of multisourced metals (sedimentary and existing mineralization) in the formation of intrusion-related stratabound deposits. Furthermore, it demonstrates the importance of integrating information regarding the source and timing of mineralization within a well-defined geological framework, which can yield information about the ore-forming processes and help to guide mineral exploration.**

## INTRODUCTION

Intrusion-related stratabound deposits are an important end member of porphyry copper systems that contribute significantly to the world's supply of copper (Cu), gold (Au), and other metals (Sillitoe, 2010). Nonmagmatic fluids in porphyry copper systems, i.e., meteoric fluids, have been ubiquitously documented (D'Errico et al., 2012; Fekete et al., 2016; Li et al., 2017a), but the current consensus is that the metals are derived from magmatic-hydrothermal systems. However, the close spatial association between metalliferous sedimentary rocks and many intrusion-related stratabound deposits raises the possibility that, in addition to magmatic-hydrothermal systems, metal-bearing strata could be a complementary origin of metals.

The expansive distribution of intrusion-related stratabound deposits in the Middle–Lower Yangtze River metallogenic belt, China (Fig. 1A), offers an excellent opportunity to examine the sources of these metals. These deposits predominantly consist of massive ores of chalcopyrite and pyrite, with gold-bearing pyrite-quartz stockworks. A unique feature of these stratabound deposits is the ubiquitous presence of colloform pyrite (Gu et al., 2007), and their close spatial association with Early Cretaceous magmatic-porphry-skarn systems (Fig. 1). Although extensively studied, the genesis of these deposits remains ambiguous, with two principal genetic models having been proposed: (1) linkage to Early Cretaceous porphyry-skarn systems (Pan and Dong, 1999; Mao et al., 2011; Pirajno and Zhou, 2015); and (2) initial formation as a late Carboniferous


seafloor exhalative formation (SEDEX) system, which was then enriched/overprinted by an Early Cretaceous magmatic-hydrothermal event (Zeng et al., 2002; Gu et al., 2007). The contrasting genetic models imply different metal enrichment mechanisms and different mineral exploration programs. For example, an Early Cretaceous magmatic-hydrothermal origin suggests that these deposits are similar to mantle-type deposits (Mao et al., 2011; Li et al., 2017b; Zhang et al., 2017) in Mexico and Chile (Meinert, 1982; Sato, 1984). Therefore, a magmatic origin of metals is most likely, with stratabound deposits being expected to center around Cretaceous granites. On the other hand, a Carboniferous SEDEX origin (Xu and Zhou, 2001; Zeng et al., 2002; Gu et al., 2007; Guo et al., 2011; Xie et al., 2014) would suggest that the metals were sourced from the underlying strata by a migrating basinal fluid, and therefore the stratabound mineralization could be more laterally extensive within the late Carboniferous limestone.

Uncertainty regarding ore genesis in the Middle–Lower Yangtze River metallogenic belt is primarily due to the lack of robust constraint on the timing of mineralization and a poor understanding of the source of the metals, which is a typical challenge for hydrothermal deposit studies. Here, based on a robust geological framework, the representative Xinqiao deposit of the Middle–Lower Yangtze River metallogenic belt was selected for a pyrite Re-Os and trace-element study, in order to provide an improved genetic understanding and yield implications for mineral exploration.

## STRATABOUND DEPOSITS OF THE MIDDLE–LOWER YANGTZE RIVER METALLOGENIC BELT

As represented by the Xinqiao Cu-S-Fe-Au deposit (eastern China), stratabound ore bodies in the Middle–Lower Yangtze River metallogenic belt are hosted primarily by limestone

\*E-mail: yang.li@yale.edu; cugliyang@126.com



**Figure 1. A: Distribution of porphyry-skarn and stratabound deposits in Middle-Lower Yangtze River metallogenic belt (China) and their spatial association with Early Cretaceous granites, modified after Pan and Dong (1999) and Mao et al. (2011). B: Geological map of Xinqiao stratabound ore deposit. C: Cross section of Xinqiao deposit and relevant chronological data. 1—Li et al. (2017b); 2—Yang et al. (2004); 3—this study; 4—Guo et al. (2011); 5—Zhang et al. (2017). Abbreviations: py—pyrite; Fm—Formation; Qtz—quartz; FI—fluid inclusion.**

and dolomite units within the late-middle Carboniferous Chuanshan-Huanglong Formations. These units lie above the early Carboniferous Gaolishan Formation sandstone (Figs. 1B and 1C). Minor massive pyritic ores are also present in overlying Permian and Triassic carbonate units. The deposits exhibit a close spatial association with Early Cretaceous (ca. 145–135 Ma) granitoids (Zhou et al., 2008; Li et al., 2010), which are hosted by Carboniferous-Permian carbonate units. Many of the granitoids are associated with porphyry and skarn mineralization (Mao et al., 2011; Pirajno and Zhou, 2015). The stratabound ore bodies predominantly consist of Cu-bearing pyrite and pyrrhotite (Li et al., 2017b). A feature unique to these deposits is pyrite exhibiting a colloform texture (Xie et al., 2014). Locally, veins bearing colloform pyrite are observed to crosscut the Permian limestone (Fig. DR3 in the GSA Data Repository<sup>1</sup>). Vertical to subvertical Au-bearing pyrite stockworks occur beneath the stratabound ore bodies and are hosted by the early Carboniferous Gaolishan Formation sandstone (Guo et al., 2011).

#### PYRITE Re-Os AND TRACE-ELEMENT RESULTS

Four styles of pyritic mineralization (Figs. 1 and 2) from Xinqiao were examined for Re-Os and trace-element analysis. Euhedral pyrite grains (py1) from the stratabound ore body have Re and <sup>192</sup>Os concentrations of 1.5–3.6 ppb and 1.7–4.0 ppt, respectively, and yield a Re-Os isochron age of 135.5 ± 4.0 Ma (Os<sub>i</sub> = 0.79 ± 0.11; n = 5; mean squared weighted deviation [MSWD] = 2.2; Fig. 2E). Colloform pyrite (py2) from the stratabound ore body possesses 1.2–9.3 ppb Re and 2.0–43.4 ppt <sup>192</sup>Os and yields a Re-Os isochron age of 136.6 ± 4.6 Ma (Os<sub>i</sub> = 1.35 ± 0.06; n = 11; MSWD = 5.4; Fig. 2E). Euhedral garnet-skarn pyrite (py3) contains 1.4–1.6 ppb Re and 1.4–2.1 ppt <sup>192</sup>Os. For py3, two analysis yielded a date of 143 ± 16 Ma and an Os<sub>i</sub> value of 0.63 ± 0.44 (Fig. 2E). The sandstone-hosted pyrites (py4) possess 57–3692 ppb Re, 1.3–43.1 ppt <sup>192</sup>Os, and negligible common Os (<0.45%; see Table DR1 in the Data Repository). The model Re-Os ages of these pyrite grains range from 173.2 ± 1.7 Ma to 378.7 ± 1.6 Ma (Fig. 2F). Overall, py1 and py3 are characterized by low trace-element abundance, but they contain moderate Mn, Cu, Pb, Zn, W, and Ag contents (Table DR2). Py2 has higher abundances of Mn, Sb, Cu, Pb, Zn, and Ag; py4 is enriched in Au, Ba, Co, Ni, and Tl (Fig. 3A). For py4, a positive correlation is observed between the total abundances of Co, Ni, Ag, Au, Tl, Ba, Re, and Re-Os model ages (Fig. 3B).

<sup>1</sup>GSA Data Repository item 2018127, deposit geology, samples, analytical methods, and pyrite Re-Os and trace-element data, is available online at <http://www.geosociety.org/datarepository/2018/> or on request from editing@geosociety.org.

## DISCUSSION AND IMPLICATIONS


### Magmatic and Sedimentary Sourced Metals for Stratabound Ore

The Re-Os ages of py1 and py2 (135.5 ± 4.0 Ma and 136.7 ± 4.6 Ma, respectively; Fig. 2E) suggest that the two types of mineralization (euhedral and colloform pyrite) were formed broadly contemporaneously and are indistinguishable from the emplacement age of the Jitou Stock (138.5 ± 1.0 Ma; Li et al., 2017b) at Xinqiao. In addition, these ages overlap with the Re-Os age of the skarn pyrite (py3, 143 ± 16 Ma; Fig. 2E). Thus, a temporal link exists between the Early Cretaceous magmatic-skarn system associated with the Jitou Stock and the stratabound mineralization. This is inconsistent with a Carboniferous SEDEX origin (Xu and Zhou, 2001; Zeng et al., 2002; Xie et al., 2014).

At the time of emplacement of the Jitou Stock, the skarn pyrite (py3) had an Os<sub>i</sub> of 0.74 ± 0.24 (Table DR1). Taking this value as the maximum estimate of the magmatic Os<sub>i</sub>, a crust-derived origin with limited mantle input (Os<sub>i</sub> = 0.13) is inferred for the Jitou Stock. This is consistent with Jitou Stock zircon depleted Hf isotope composition (ε<sub>Hf</sub> = ~-11; Zhang et al., 2017). The similar Os<sub>i</sub> values (0.79 ± 0.11 vs. 0.74 ± 0.24) for py1 and py3, respectively, imply that the Os, and by inference the associated metals, was predominantly sourced from the Early Cretaceous magmatic-hydrothermal system.

It has been previously proposed that the colloform pyrite was initially formed in the Carboniferous (Xu and Zhou, 2001; Zeng et al., 2002; Gu et al., 2007; Xie et al., 2014) and then recrystallized to euhedral pyrite during the Early Cretaceous magmatic-hydrothermal event. This scenario is not supported by the sharp contact relationship between the euhedral and colloform pyrites (Fig. DR2A), nor by the crosscutting relationship between the colloform pyrite and Permian limestones at Dongguashan and Wushan (Fig. DR3). In addition, the Os<sub>i</sub> (Fig. 2E) and trace-element abundances (Figs. 3A and 3B) of py2 are distinct from py1; hence, a recrystallization origin of py1 from py2 is unlikely. Further, the colloform pyrite is composed of fine-grained (80 nm to 1.5 μm) cubic crystals and not framboids, which is inconsistent with a sedimentary origin (Sweeney and Kaplan, 1973).

The highly radiogenic Os<sub>i</sub> value (1.35 ± 0.06) of the colloform pyrite indicates that the Os, and by inference the other metals, was not solely magmatically derived. In the Xinqiao area, the most likely source to provide a radiogenic <sup>187</sup>Os/<sup>188</sup>Os composition is the Late Permian metalliferous black shales. These shales are enriched in Re (403–1002 ppb) and Os (0.3–1 ppb) and yield an Early Cretaceous <sup>187</sup>Os/<sup>188</sup>Os composition of 7.6 ± 3.8 (Yang et al., 2004). Therefore, the most geological plausible scenario is that the colloform pyrite was formed through intensive water-rock



**Figure 2. Representative pyrite mineralization at Xinqiao (eastern China) and Re-Os ages.** A: Euhedral pyrite grains (py1) cemented by calcite from stratabound ore body. B: Colloform pyrite (py2) from stratabound ore body, which is composed of fine-grained (80 nm to 1.5  $\mu$ m) cubic pyrite grains, with local distribution of calcite grains. No framboidal pyrite grains were observed in this study. C: Pyrite samples (py3) from garnet-bearing skarn ore. D: Pyrite samples (py4) hosted by sandstone beneath stratabound ore body. E: Re-Os isochrons of py1, py2, and py3. F: Model ages of py4 and their correlation with Re abundances. Also shown are data from Guo et al. (2011). Resin refers to materials used for making mounts. Py—pyrite, cal—calcite, qtz—quartz, gar—garnet, MSWD—mean squared weighted deviation. All uncertainties are at 2 $\sigma$  level.

interaction between the Early Cretaceous magmatic-hydrothermal fluids and the Late Permian metalliferous black shales; this is further supported by the trace element data (Fig. 3A).

### Recycling Existing Mineralization for the Au-Bearing Pyrite Stockworks


The Au-bearing pyrite (py4) veins hosted by the sandstone underlying the stratabound ore body are dated at  $138 \pm 2.3$  Ma (quartz fluid inclusion Rb-Sr; Zhang et al., 2017), suggesting a temporal and genetic association with the Jitou Stock, which is further supported by fluid inclusion and  $\delta^{18}\text{O}$  data from quartz coprecipitated with py4 (up to 597  $^{\circ}\text{C}$  and 63.7% NaCl equiv.;  $6.81\text{‰} \pm 2.76\text{‰} \delta^{18}\text{O}$ ; Wang and Ni, 2009; Li et al., 2017b). In contrast, a  $319 \pm 13$  Ma Re-Os date ( $n = 9$ ; MSWD = 13; Guo et al., 2011) for py4 seems consistent with the hypothesis that these pyrite veins were the fluid conduit (stockwork feeder) for a Carboniferous SEDEX system (Xu and Zhou, 2001; Zeng et al., 2002; Gu et al., 2007). However, a SEDEX scenario is not supported by the following observations. First, py4 Re-Os data from both this study and Guo et al. (2011) share similar characteristics (enriched in Re and limited to no common Os) and yield highly variable model ages (Fig. 2F; 379–173 Ma). Therefore, the py4 Re-Os data do not meet the necessary criteria for isochron dating. Second, py4 does not contain common Os, but in SEDEX systems, the basal fluid must interact with the basement rocks, and hence should inherit common Os with a radiogenic  $\text{Os}_i$  value (Hnatyshin et al., 2015). Third, the pyrite veins beneath the stratabound ore body are structurally controlled and only possess silicified and sericite-bearing selvages (Wang and Ni, 2009; Guo et al., 2011; Li et al., 2017b). In contrast, the fluid conduits in SEDEX systems are developed in synsedimentary

faults and are characterized by tourmaline- and albite-bearing alteration assemblages (Leach et al., 2005). As such, in accordance with the  $138 \pm 2.3$  Ma Rb-Sr age (Zhang et al., 2017), we suggest that the sandstone-hosted mineralization was temporally associated with the Early Cretaceous magmatic-hydrothermal system.

For py4, in order to yield geologically reasonable ages (ca. 138 Ma), each sample had to be corrected using widely different and geologically implausible  $\text{Os}_i$  values (e.g., –223 to 59; Table DR1). Given the fact that these samples possess very high abundances of Re and radiogenic Os with negligible common Os, it is unlikely that the observed ages were caused by disturbance of the Re-Os systematics. In this regard, the most plausible genesis for py4 is the leaching of rocks enriched in Re and radiogenic Os by the Early Cretaceous magmatic-hydrothermal fluid. A higher degree of water-rock interaction resulted in the inheritance of more Re,  $^{187}\text{Os}$ , and trace elements, consistent with the positive correlations between model ages of py4 and their Re and trace-element concentrations (Figs. 2F and 3). Rhenium and Os concentrations are typically very low in crustal rocks (0.20 and 0.03 ppb, respectively), but they are known to be high in molybdenite (e.g., hundreds to thousands of ppm for Re, and up to ppm levels for  $^{187}\text{Os}$ ) and to a lesser extent in other sulfides (Selby et al., 2009; Stein, 2014). As such, the most likely source for the elevated Re (ppm level), radiogenic Os ( $^{187}\text{Os}$ , ppb level), and negligible common Os in py4 is porphyry-style Au-Mo mineralization.

### Implications for Ore Genesis and Origin of the Metals

In the Middle–Lower Yangtze River metallogenic belt, stratabound deposits are all spatially




**Figure 3. Trace-element compositions of the studied pyrite samples.** A: Py2 is characterized by high total abundances of Co, Ni, Ag, Au, Tl, and Ba; py4 is enriched in Cu, W, Pb, and Zn. In contrast, py1 and py3 are depleted in these trace elements. B: Model ages of py4 are positively correlated with total abundances of Co, Ni, Ag, Au, Tl, and Ba. Uncertainties are smaller than symbol size.

associated with the Early Cretaceous magmatic-porphyry-skarn systems (Fig. 1A), and the ore-forming fluids and alteration assemblages have a close magmatic affinity, but they show few characteristics of typical SEDEX systems (Pan



and Dong, 1999; Mao et al., 2011). As such, an Early Cretaceous intrusion-related carbonate replacement origin is suggested, with the stratabound ore bodies being spatially controlled by the unconformity between the early Carboniferous sandstone and late Carboniferous carbonate units, and the Au-bearing stockworks being a product of recycled preexisting porphyry-style mineralization by the Early Cretaceous magmatic-hydrothermal fluid (Fig. 4).



**Figure 4. Sketch illustrating genesis of intrusion-related stratabound deposits in Middle–Lower Yangtze metallogenic belt (China), which are temporally and spatially associated with Early Cretaceous granites, porphyry, and skarn deposits. Model highlights the importance of recycling metals from metalliferous sedimentary rocks and preexisting mineralization in formation of intrusion-related stratabound deposits; py—pyrite.**

This study highlights the benefits of coupling absolute timing and source constraints for mineralization with a detailed geological framework, which can significantly advance our understanding of the ore-forming process of intrusion-related stratabound deposits, and underpin exploration models. Further, the recycling of metals from metalliferous sedimentary rocks (i.e., black shales) and preexisting mineralization by magmatic-hydrothermal fluids could be important mechanisms in ore formation.

#### ACKNOWLEDGMENTS

Y. Li acknowledges a two-month fellowship and a grant (SKL-K201706) from the Institute of Geology and Geophysics, Chinese Academy of Sciences, and extends his gratitude to Jian-Wei Li for his encouragement during this study. Selby acknowledges the Total Endowment Fund and Dida Scholarship from the China University of Geosciences, Wuhan. We thank Zhen-Ting Jiang for assistance with scanning electron microscope imaging, and Ann Bauer, Alan Rooney, and Rachael Bullock for editorial comments, and we extend our thanks to the editor and reviewers of *Geology* for their insightful comments.

#### REFERENCES CITED

D’Errico, M.E., Lackey, J.S., Surpless, B.E., Loewy, S.L., Wooden, J.L., Barnes, J.D., Strickland, A., and Valley, J.W., 2012, A detailed record of shallow hydrothermal fluid flow in the Sierra Nevada magmatic arc from low- $\delta^{18}\text{O}$  skarn garnets:

*Geology*, v. 40, p. 763–766, <https://doi.org/10.1130/G33008.1>.

Fekete, S., Weis, P., Driesner, T., Bouvier, A.S., Baumgartner, L., and Heinrich, C.A., 2016, Contrasting hydrological processes of meteoric water incursion during magmatic-hydrothermal ore deposition: An oxygen isotope study by ion microprobe: *Earth and Planetary Science Letters*, v. 451, p. 263–271, <https://doi.org/10.1016/j.epsl.2016.07.009>.

Gu, L.X., Zaw, K., Hu, W.X., Zhang, K.J., Ni, P., He, J.X., Xu, Y.T., Lu, J.J., and Lin, C.M., 2007, Distinctive features of late Palaeozoic massive sulphide deposits in South China: *Ore Geology Reviews*, v. 31, p. 107–138, <https://doi.org/10.1016/j.oregeorev.2005.01.002>.

Guo, W.M., Lu, J.J., Jiang, S.Y., Zhang, R.Q., and Qi, L., 2011, Re-Os isotope dating of pyrite from the foot-wall mineralization zone of the Xinqiao deposit, Tongling, Anhui Province: Geochronological evidence for submarine exhalative sedimentation: *Chinese Science Bulletin*, v. 56, p. 3860–3865, <https://doi.org/10.1007/s11434-011-4770-y>.

Hnatyshin, D., Creaser, R.A., Wilkinson, J.J., and Gleeson, S.A., 2015, Re-Os dating of pyrite confirms an early diagenetic onset and extended duration of mineralization in the Irish Zn-Pb ore field: *Geology*, v. 43, p. 143–146, <https://doi.org/10.1130/G36296.1>.

Leach, D., Sangster, D., Kelley, K., Large, R., Garven, G., Allen, C., Gutzmer, J., and Walters, S.G., 2005, Sediment-hosted lead-zinc deposits: A global perspective: *Economic Geology and the Bulletin of the Society of Economic Geologists*, v. 100, p. 561–607.

Li, X.H., Li, W.X., Wang, X.C., Li, Q.L., Liu, Y., Tang, G.Q., Gao, Y.Y., and Wu, F.Y., 2010, SIMS U-Pb zircon geochronology of porphyry Cu-Au-(Mo) deposits in the Yangtze River metallogenic belt, eastern China: Magmatic response to Early Cretaceous lithospheric extension: *Lithos*, v. 119, p. 427–438, <https://doi.org/10.1016/j.lithos.2010.07.018>.

Li, Y., Li, X.H., Selby, D., and Li, J.W., 2017a, Pulsed magmatic fluid release for the formation of porphyry deposits: Tracing fluid evolution in absolute-time from the Tibetan Qulong Cu-Mo deposit: *Geology*, v. 46, p. 7–11, <https://doi.org/10.1130/G39504.1>.

Li, Y., Li, J.W., Li, X.H., Selby, D., Huang, G.H., Chen, L.J., and Zheng, K., 2017b, An Early Cretaceous carbonate replacement origin for the Xinqiao stratabound massive sulfide deposit, Middle–Lower Yangtze metallogenic belt, China: *Ore Geology Reviews*, v. 80, p. 985–1003, <https://doi.org/10.1016/j.oregeorev.2016.08.017>.

Mao, J.W., Xie, G.Q., Duan, C., Pirajno, F., Ishiyama, D., and Chen, Y.C., 2011, A tectono-genetic model for porphyry-skarn-stratabound Cu-Au-Mo-Fe and magnetite-apatite deposits along the Middle–Lower Yangtze River Valley, eastern China: *Ore Geology Reviews*, v. 43, p. 294–314, <https://doi.org/10.1016/j.oregeorev.2011.07.010>.

Meinert, L.D., 1982, Skarn, manto, and breccia pipe formation in sedimentary-rocks of the Cananea Mining District, Sonora, Mexico: *Economic Geology and the Bulletin of the Society of Economic Geologists*, v. 77, p. 919–949, <https://doi.org/10.2113/gsecongeo.77.4.919>.

Pan, Y.M., and Dong, P., 1999, The Lower Changjiang (Yangzi/Yangtze River) metallogenic belt, east central China: Intrusion- and wall rock-hosted Cu-Fe-Au, Mo, Zn, Pb, Ag deposits: *Ore Geology Reviews*, v. 15, p. 177–242, [https://doi.org/10.1016/S0169-1368\(99\)00022-0](https://doi.org/10.1016/S0169-1368(99)00022-0).

Pirajno, F., and Zhou, T., 2015, Intracratonic porphyry and porphyry-skarn mineral systems in eastern China: Scrutiny of a special case “Made-in-China”: *Economic Geology and the Bulletin of the Society of Economic Geologists*, v. 110, p. 603–629, <https://doi.org/10.2113/econgeo.110.3.603>.

Sato, T., 1984, Manto type copper deposits in Chile: A review: *Bulletin of the Geological Survey of Japan*, v. 35, p. 565–582.

Selby, D., Kelley, K.D., Hitzman, M.W., and Zieg, J., 2009, Re-Os sulfide (bornite, chalcopyrite, and pyrite) systematics of the carbonate-hosted copper deposits at Ruby Creek, southern Brooks Range, Alaska: *Economic Geology and the Bulletin of the Society of Economic Geologists*, v. 104, p. 437–444, <https://doi.org/10.2113/gsecongeo.104.3.437>.

Sillitoe, R.H., 2010, Porphyry copper systems: *Economic Geology and the Bulletin of the Society of Economic Geologists*, v. 105, p. 3–41, <https://doi.org/10.2113/gsecongeo.105.1.3>.

Stein, H.J., 2014, Dating and tracing the history of ore formation, in Turekian, H.D.H.K., ed., *Treatise on Geochemistry* (2nd ed.): Oxford, UK, Elsevier, p. 87–118, <https://doi.org/10.1016/B978-0-08-095975-7.01104-9>.

Sweeney, R.E., and Kaplan, I.R., 1973, Pyrite framboid formation: Laboratory synthesis and marine sediments: *Economic Geology*, v. 68, p. 618–634, <https://doi.org/10.2113/gsecongeo.68.5.618>.

Wang, G.-G., and Ni, P., 2009, Fluid inclusions study for the pyrite vein beneath Xinqiao stratiform massive Cu deposit: *Acta Mineralogica Sinica*, v. 29, p. 256 [in Chinese].

Xie, Q.-Q., Chen, T.-H., Fan, Z.-L., Xu, X.-C., Zhou, Y.-F., Shi, W.-B., and Xue, J.-J., 2014, Morphological characteristics and genesis of colloform pyrite in Xinqiao Fe-S deposit, Tongling, Anhui Province: *Scientia Sinica Terrae*, v. 44, p. 2665–2674 [in Chinese].

Xu, G., and Zhou, J., 2001, The Xinqiao Cu-S-Fe-Au deposit in the Tongling mineral district, China: Synorogenic remobilization of a stratiform sulfide deposit: *Ore Geology Reviews*, v. 18, p. 77–94, [https://doi.org/10.1016/S0169-1368\(01\)00017-8](https://doi.org/10.1016/S0169-1368(01)00017-8).

Yang, G., Chen, J.F., Du, A.D., Qu, W.J., and Yu, G., 2004, Re-Os dating of the Mo-bearing black shale from Laoyalong, Anhui: *Chinese Science Bulletin*, v. 49, p. 1205–1208 [in Chinese].

Zeng, P.-S., Pei, R.-F., and Hou, Z.-Q., 2002, Sedex-type massive sulfide deposits in Tongling block, Anhui: *Mineralium Deposita*, v. 21, p. 532–535 [in Chinese].

Zhang, Y., Shao, Y.J., Li, H.B., and Liu, Z.F., 2017, Genesis of the Xinqiao Cu-S-Fe-Au deposit in the Middle–Lower Yangtze River Valley metallogenic belt, eastern China: Constraints from U-Pb-Hf, Rb-Sr, S, and Pb isotopes: *Ore Geology Reviews*, v. 86, p. 100–116, <https://doi.org/10.1016/j.oregeorev.2017.02.014>.

Zhou, T.F., Yu, F., and Yuan, F., 2008, Advances on petrogenesis and metallogeny study of the mineralization belt of the Middle and Lower Reaches of the Yangtze River area: *Yanshi Xuebao*, v. 24, p. 1665–1678 [in Chinese with English abstract].

Manuscript received 14 December 2017

Revised manuscript received 30 January 2018

Manuscript accepted 31 January 2018

Printed in USA