

Contributions to Geology

The University of Wyoming

Volume 18, Number 2

Spring, 1980

CONTENTS

	<i>Page</i>	
<i>P. S. Dahl and L. M. Friberg</i>	77	The occurrence and chemistry of epidote-clinozoisites in mafic gneisses from the Ruby Range, southwestern Montana
<i>D. L. Blackstone, Jr.</i>	83	Foreland deformation: compression as a cause
<i>John A. Dorr, Jr. and Philip D. Gingerich</i>	101	Early Cenozoic mammalian paleontology, geologic structure, and tectonic history in the overthrust belt near LaBarge, western Wyoming
<i>Carter Lupton, Diane Gabriel, and Robert M. West</i>	117	Paleobiology and depositional setting of a Late Cretaceous vertebrate locality, Hell Creek Formation, McCone County, Montana
<i>Michael E. Nelson, James H. Madsen, Jr., and W. Lee Stokes</i>	127	A titanotheres from the Green River Formation, central Utah: <i>Teleodus uintensis</i> (Perissodactyla: Bronthoheriidae)
<i>Bruce J. MacFadden</i>	135	Eocene perissodactyls from the type section of the Tepee Trail Formation of northwestern Wyoming

Cover Illustration:

View to the southwest across the valley of Wagonhound Creek to the east face of Elk Mountain, Carbon County, Wyoming. The Elk Mountain structure is asymmetrical to the east, and the roll over of sediments is visible at the extreme right center. Photograph by D. L. Blackstone, Jr. (see article by D. L. Blackstone, Jr.)

SUBSCRIPTIONS and related matters may be addressed to Circulation Dept., Contributions to Geology, P.O. Box 3006, University Station, Laramie, WY 82071, U.S.A.

POTENTIAL CONTRIBUTORS should consult inside front cover.